

## RAPID KNOCKDOWN CONTROL OF UNWANTED WEEDS

**PMRA REG. NUMBER**  
35229

**ACTIVE INGREDIENT**  
Glufosinate-ammonium  
150 g/L

**FORMULATION**  
Solution

**HRAC NUMBER**  
10

**SIGNAL WORD**  
Warning Poison

**PACKAGE SIZE**  
500 L tote

**RESTRICTED USE**  
No

Flamora™ 150 SN provides control of grassy and broadleaf weeds in canola and soybean varieties or hybrids that are specially developed to be tolerant to glufosinate ammonium.

- Works as a contact herbicide to deliver nonselective control of weeds
- Excellent rotational product for herbicide resistance management
- Tank-mix compatibility with select herbicides for enhanced activity

### KEY CROPS



CANOLA



SOYBEANS

### KEY WEEDS CONTROLLED



TARGET  
GRASSY &  
BROADLEAF

- Barnyard Grass
- Canada Thistle
- Cleavers
- Common Chickweed
- Cow Cockle
- Dandelion
- Flixweed
- Green Foxtail
- Hemp-nettle
- Kochia
- Lady's-thumb
- Lamb's-quarters
- Perennial Sow Thistle
- Pigweed
- Quackgrass
- Redroot
- Round-leaved Mallow
- Russian Thistle
- Scentless Chamomile
- Shepherd's-purse
- Smartweed
- Stinkweed
- Stork's-bill
- Volunteer Barley
- Volunteer Flax
- Volunteer Wheat
- Wild Buckwheat
- Wild Oats
- Wild Mustard

### WATER VOLUME



**GROUND APPLICATION:** 10 GAL/AC or 45.5 L/AC

**AERIAL APPLICATION:** 6 GAL/AC or 27.3 L/AC





## RATES FOR KEY WEEDS CONTROLLED

### 1.33 L/HA = 0.53 L/AC

Cow Cockle	
Green Foxtail	Maximum 3 Tillers

### 2 L/HA = 0.81 L/AC

Barnyard Grass	
Lady's Thumb	
Lamb's-quarters	
Russian Thistle	Up to 8 cm height
Smartweed	
Stinkweed	
Volunteer Flax	Up to 6 cm height
Wild Mustard	

### 2.67 L/HA = 1.08 L/AC

Canada Thistle <sup>1</sup>	Up to 10 cm height
Common Chickweed	Leaf Pairs
Hemp-nettle	Leaf Pairs
Kochia	Up to 8 cm height
Perennial Sow Thistle	
Quackgrass <sup>1</sup>	
Redroot	
Pigweed	
Round-leaved Mallow	
Scentless Chamomile	Up to 10 cm height
Shepherd's-purse	
Volunteer Barley <sup>2</sup>	Maximum 2 Tillers
Volunteer Wheat	Maximum 2 Tillers
Wild Buckwheat	

### 3.33 L/HA = 1.35 L/AC

Cleavers	Whorls
Dandelion	1-15 cm rosette
Flixweed	Up to 10 cm height
Hemp-nettle	Leaf Pairs
Stork's-bill	
Wild Oats	Maximum 2 Tillers
Canada Thistle <sup>1</sup>	Up to 10 cm height
Quackgrass <sup>1</sup>	
Volunteer Barley <sup>2</sup>	Maximum 2 Tillers
Volunteer Wheat	Maximum 2 Tillers
Wild Buckwheat	

### 4 L/HA = 1.62 L/AC

Canada Thistle <sup>3</sup>	Up to 10 cm height
Quackgrass <sup>4</sup>	

<sup>1</sup>Top Growth Suppression Only

<sup>2</sup>Suppression Only

<sup>3</sup>Better Top Growth Suppression

<sup>4</sup>Season-long Control for Heavy Populations

## APPLICATION TIMING: CANOLA

Staging for first or second application: cotyledon until prior to bolting.

A second application of Flamora 150 SN can be made to canola treated initially with up to 4 L/ha (1.62 L/ac) if new weed germination or growth is present. This first application of up to 4 L/ha (1.62 L/ac) may be followed by a second application of up to 3.33 L/ha (1.35 L/ac). Do not apply more than a total of 7.33 L/ha (2.97 L/ac) per year.

- **Rainfastness:** If rainfall occurs within 4 hours of application, effectiveness may be reduced
- **Tankmixes and Mixing Order:** Talk to your retailer agronomist for recommendation and guidance