

MONFERNO™

250 EC

FUNGICIDE

PROTECT THE YIELD. POWER THE CROP.

PMRA REG. NUMBER
35775

ACTIVE INGREDIENT
Pyraclostrobin 250 g/L

FORMULATION
Emulsifiable
Concentrate

FRAC NUMBER
11

SIGNAL WORD
DANGER: POISON

RE-ENTRY INTERVAL
12 hours

PACKAGE SIZE
2 x 6.5 L jugs/case
500 L tote

RESTRICTED USE
No

Monferno™ 250 EC delivers reliable, broad-spectrum protection powered by pyraclostrobin. It promotes healthier plants with improved stress tolerance, strong leaf retention, and yield preservation from early growth through harvest.

- Broad-spectrum control of major foliar diseases
- Dual preventive and curative activity
- Enhances plant health and stress tolerance
- Provides translaminar protection for both sides of the leaf

KEY CROPS



CEREALS



CANOLA



FLAX



SOYBEANS



PEAS



LENTILS +
CHICKPEAS



POTATOES

KEY DISEASES CONTROLLED OR SUPPRESSED



TARGET
DISEASES

- Anthracnose
- Alternaria leaf spot
- Ascochyta blight
- Black spot
- Blackleg
- Cercospora leaf spot
- Crown rust
- Downy mildew
- Early and late blight
- Leaf rust
- Mycosphaerella blight
- Net blotch
- Pasmó
- Powdery mildew
- Scald
- Septoria leaf blotch
- Spot blotch
- Stripe rust
- Tan spot

WATER VOLUME



GROUND: Minimum 10 GAL/AC or 45.5 L/AC

AERIAL: Minimum 5 GAL/AC or 22.75 L/AC



AVESTACROP.COM



1-800-607-5443



INFO@AVESTACROP.COM



Avesta™
CropScience

APPLICATION INFORMATION

Monferno™ 250 EC can be applied before or after infection, but the best results come from a proactive approach. Applying Monferno preventively as part of a regular spray program—and rotating with fungicides from other groups—helps ensure strong, consistent disease control throughout the season.

Apply Monferno 250 EC after flag leaf emergence (GS 37–39) to protect the flag leaf and support maximum yield potential. Use the higher rate when disease pressure is high or conditions favor infection.

If disease persists, apply again 10–14 days later using a fungicide with a different mode of action.

CROP	DISEASE	APPLICATION RATE* (L/AC)
WHEAT	Leaf rust Tan spot Septoria leaf spot	0.121-0.243
	Spot blotch Stripe rust Powdery mildew	0.162-0.243
BARLEY	Net blotch	0.121-0.243
	Spot blotch Stripe rust Scald	0.162-0.243
RYE	Leaf rust	0.121-0.243
	Powdery mildew	0.126-0.243
OATS	Crown rust	0.121-0.162
CANOLA	Black spot (<i>Alternaria brassicae</i> and <i>Alternaria raphani</i>) Blackleg (<i>Leptosphaeria maculans</i>)	0.121-0.162
FLAX	Pasmo (<i>Septoria linicola</i>)	0.121-0.162
SOYBEANS	Frog eye leaf spot (<i>Cercospora sojina</i>) Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	0.162-0.243
PEAS	Mycosphaerella blight (<i>Mycosphaerella</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.)	0.162
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	0.162-0.243
LENTILS	Anthrachnose (<i>Colletotrichum</i> spp.) Ascochyta blight (<i>Ascochyta</i> spp.)	0.162
CHICKPEAS	Ascochyta blight (<i>Ascochyta rabiei</i>)	Monferno250EC must be tank mixed at a rate of 0.162-0.243 L/ac with LANCE® WDG Fungicide at 0.146 – 0.170 kg/ac.
POTATOES	Early blight (<i>Alternaria solani</i>)	0.182-0.217
	Late blight (<i>Phytophthora infestans</i>)	0.182-0.217

* Do not apply more than 0.162 L/ac by aerial application.