STRYDOR™ 360



HERBICIDE

# CONTROL THISTLE – CLEAR FIELDS

**PMRA REG. NUMBER**
36160

**ACTIVE INGREDIENT**
Clopyralid 360 g/L

**FORMULATION**
Soluble Liquid

**HRAC NUMBER**
4

**SIGNAL WORD**
WARNING: POISON

**RE-ENTRY INTERVAL**
12 hours

**PACKAGE SIZE**
4 × 4.45 L jugs/case

**RESTRICTED USE**
No

Strydor™ 360 delivers powerful, lasting control of tough weeds like Canada thistle and sow thistle in canola, cereals, and pastures. Its systemic activity provides extended residual performance and reliable, season-long results.

- Controls annual and perennial broadleaf weeds
- Moves throughout the entire plant system for lasting control
- Provides extended residual performance
- Flexible, easy-to-use formulation

## KEY CROPS

CANOLA

CEREALS

PASTURES

## KEY WEEDS CONTROLLED

TARGET BROADLEAF WEEDS

- Alsike clover
- Canada thistle
- Common groundsel
- Common ragweed
- Oxeye daisy*
- Perennial sow-thistle*
- Scentless chamomile
- Sheep sorrel*
- Vetch
- Volunteer alfalfa
- Wild buckwheat

**Suppression Only*

## WATER VOLUME

**GROUND:** 10–20 GAL/AC or 45.5–91 L/AC (Typical)

AVESTACROP.COM
1-800-607-5443
INFO@AVESTACROP.COM
LinkedIn
Avesta™ CropScience

STRYDOR™ 360



## APPLICATION INFORMATION

| CROP | RATE (L/AC) | APPLICATION METHOD | APPLICATION VOLUME (L/AC) | CROP TIMING | NOTES |
|---|---|---|---|---|---|
| **WHEAT** SPRING | 0.136–0.204 | Ground | 40.5–80.9 | 3-leaf → early flag leaf | Use lower rate for 6–8 weeks control or higher rate for season-long thistle control. |
| **BARLEY** SPRING | 0.136–0.204 | Ground | 40.5–80.9 | 3-leaf to flag-leaf | Do not undersow legumes. |
| **OATS** | 0.136–0.204 | Ground | 40.5–80.9 | 3-leaf to flag-leaf | See Grazing and Harvesting for Feed Section of label for intervals. |
| **CANOLA** WESTERN CANADA | 0.204–0.403 | Ground | 40.5–80.9 | 2–6 leaf stage | For use on Polish and Argentine varieties. |
| **FLAX** | 0.272–0.403 | Ground | 40.5–80.9 | 5–10 cm crop height | Higher rate can extend thistle control into following year. |
| **PASTURE & RANGELAND** | 0.204–0.403 | Ground | 44.5–89 | Active weed growth | Apply once per season; for established grasses, apply at shot-blade stage or early spring. |

## ADDITIONAL INFORMATION

- Commonly used in tank-mixes with MCPA, 2,4-D, or glyphosate to expand control of weeds.
- Add Strydor™ 360 first to the spray tank, followed by emulsifiable concentrates, and then adjuvants.
- Maintain continuous agitation throughout mixing and spraying.
- Do not apply by air.
- Follow the most restrictive partner label for rates, timing, and buffer zones.

**AVESTACROP.COM**

*Refer to product label for complete application and mixing instructions.*

©2025 Avesta CropScience Inc. Important: Always read and follow label instructions. Not all products are registered for use in every province. Strydor™ is a trademark of Avesta CropScience Inc.

Avesta™ CropScience